

# Gamma Glutamyl Transferase (GGT)

CPT: 82977

## Summary of CMS National Coverage Policies\*

### Coverage Indications, Limitations, and/or Medical Necessity

Gamma glutamyl transferase (GGT) is an intracellular enzyme that appears in blood following leakage from cells. Renal tubules, liver, and pancreas contain high amounts, although the measurement of GGT in serum is almost always used for assessment of Hepatobiliary function. Unlike other enzymes which are found in heart, skeletal muscle, and intestinal mucosa as well as liver, the appearance of an elevated level of GGT in serum is almost always the result of liver disease or injury.

It is specifically useful to differentiate elevated alkaline phosphatase levels when the source of the alkaline phosphatase increase (bone, liver, or placenta) is unclear. The combination of high alkaline phosphatase and a normal GGT does not, however, rule out liver disease completely. As well as being a very specific marker of Hepatobiliary function, GGT is also a very sensitive marker for hepatocellular damage. Abnormal concentrations typically appear before elevations of other liver enzymes or biliuria are evident. Obstruction of the biliary tract, viral infection (e.g., hepatitis, mononucleosis), metastatic cancer, exposure to hepatotoxins (e.g., organic solvents, drugs, alcohol), and use of drugs that induce microsomal enzymes in the liver (e.g., cimetidine, barbiturates, phenytoin, and carbamazepine) all can cause a moderate to marked increase in GGT serum concentration. In addition, some drugs can cause or exacerbate liver dysfunction (e.g., atorvastatin, troglitazone, and others as noted in FDA Contraindications and Warnings.)

GGT is useful for diagnosis of liver disease or injury, exclusion of hepatobiliary involvement related to other diseases, and patient management during the resolution of existing disease or following injury.

### Indications

1. To provide information about known or suspected hepatobiliary disease, for example:
  - a. Following chronic alcohol or drug ingestion
  - b. Following exposure to hepatotoxins
  - c. When using medication known to have a potential for causing liver toxicity (e.g., following the drug manufacturer's recommendations)
  - d. Following infection (e.g., viral hepatitis and other specific infections such as amebiasis, tuberculosis, psittacosis, and similar infections)
2. To assess liver injury/function following diagnosis of primary or secondary malignant neoplasms
3. To assess liver injury/function in a wide variety of disorders and diseases known to cause liver involvement (e.g., diabetes mellitus, malnutrition, disorders of iron and mineral metabolism, sarcoidosis, amyloidosis, lupus, and hypertension)
4. To assess liver function related to gastrointestinal disease
5. To assess liver function related to pancreatic disease
6. To assess liver function in patients subsequent to liver transplantation
7. To differentiate between the different sources of elevated alkaline phosphatase activity

### Limitations

When used to assess liver dysfunction secondary to existing non-hepatobiliary disease with no change in signs, symptoms, or treatment, it is generally not necessary to repeat a GGT determination after a normal result has been obtained unless new indications are present. If the GGT is the only "liver" enzyme abnormally high, it is generally not necessary to pursue further evaluation for liver disease for this specific indication.

When used to determine if other abnormal enzyme tests reflect liver abnormality rather than other tissue, it generally is not <sup>necessary</sup> to repeat a GGT more than one time per week.

Because of the extreme sensitivity of GGT as a marker for cytochrome oxidase induction or cell membrane permeability, it is generally not useful in monitoring patients with known liver disease.

## Gamma Glutamyl Transferase (GGT)

CPT: 82977

The ICD10 codes listed below are the top diagnosis codes currently utilized by ordering physicians for the limited coverage test highlighted above that are also listed as medically supportive under Medicare's limited coverage policy. If you are ordering this test for diagnostic reasons that are not covered under Medicare policy, an Advance Beneficiary Notice form is required.

\*Please refer to the Limitations or Utilization Guidelines section on previous page(s).

CODE	DESCRIPTION
C61	Malignant neoplasm of prostate
E11.22	Type 2 diabetes mellitus with diabetic chronic kidney disease
E11.65	Type 2 diabetes mellitus with hyperglycemia
E11.9	Type 2 diabetes mellitus without complications
E78.00	Pure hypercholesterolemia, unspecified
E78.2	Mixed hyperlipidemia
E78.49	Other hyperlipidemia
E78.5	Hyperlipidemia, unspecified
E78.9	Disorder of lipoprotein metabolism, unspecified
E83.40	Disorders of magnesium metabolism, unspecified
E83.42	Hypomagnesemia
K74.60	Unspecified cirrhosis of liver
K75.81	Nonalcoholic steatohepatitis (NASH)
K76.0	Fatty (change of) liver, not elsewhere classified
K76.89	Other specified diseases of liver
K76.9	Liver disease, unspecified
R74.01	Elevation of levels of liver transaminase levels
R74.8	Abnormal levels of other serum enzymes
Z79.899	Other long term (current) drug therapy
Z94.4	Liver transplant status

To view the complete policy and the full list of codes, please refer to the CMS website [Home - Centers for Medicare & Medicaid Services | CMS](#)